



Ecological Data



- Data comes from many sources
 - Multiple entities fly satellites and operate fixed sensors – imagery from NASA, precipitation from NOAA, discharge from USGS, water quality from EPA
 - Scientists take data in field campaigns, sensor or other instrument deployments, photographs
 - Collecting site properties such as channel cross-section or leaf area index done by individuals as well as agencies
- Data repositories not universal
 - passed around in e-mail
 - edited at will
 - silently corrected in repositories when "everybody knows"
- Data comes from unexpected places
 - Published literature supplies various numbers and graphs giving context, sanity checks on the results
- Analysis routines usually custom



Example Dataset - FLUXNET




- Global carbon flux and micrometeorological measurements
 - 960 site years of data
 - 40 variables and 60 quality tags
 - 1/2 hourly measurements
 - 253 sites
 - ~120 site investigators (collecting data)
 - 72 synthesis paper proposals involving ~110 collaborators
- Goal: Understand the role of photosynthesis in carbon cycle



Windows HPC Server 2008

Developing Common Infrastructure

- Data ingested into a database
- Database schema organized by dataset, site, time, and variable
- On-line analytical processing (OLAP) data cubes provide browsing access
- Reports provide web pages describing the data
- Ancillary data gathered and curated
- Collaboration tools facilitate communication between data providers and data users as well as between data users



Interaction and Visualization



- Mashup maps
 - locate sensors
- Data summary reports
 - Data browsing
 - Data selection
 - Cross-site information
- OLAP data cube interfaces
 - Matlab analysis and plotting
 - Excel data browsing and plotting
- Collaboration capabilities
 - Improve communication between data producers and data consumers



Leveraging a Common Approach



and Technology Base

Established Projects

- Russian River
 - Digital watershed built from existing data to look at changes over time affecting salmon
- Carbon Flux Measurement Network
 - Regional (AmeriFlux) and global (FLUXNET) datasets compiled from local carbon flux measurements
- Pajaro River
 - Digital watershed built from existing data to look at natural channel flood control alternatives (listed in top 10 "endangered river" in 2006)

Projects Getting Underway Now

- National Marine Fisheries Service
 - Digital watershed to support fish recovery planning for northern California
- California watersheds server
- North American Carbon Program Carbon flux modeling

Acknowledgements



Berkeley Water Center, University of

California, Berkeley, Lawrence

Berkeley Laboratory

Jim Hunt (UCB)

Deb Agarwal (LBL)

Robin Weber (UCB)

Monte Good (LBL)

Rebecca Leonardson (UCB student)

Nick Butler (UCB student)

Matt Rodriguez (SDSC Student)

Carolyn Remick (UCB) Susan Hubbard (LBL)

University of Virginia

Marty Humphrey Norm Beekwilder

Microsoft

Catharine van Ingen

Bora Beran

Jayant Gupchup (student)

Savas Parastidis

Andy Sterland

Nolan Li (student)

Tony Hey

Dan Fay

Jing De Jong-Chen

Stuart Ozer

SQL product team

Jim Gray

Ameriflux Collaboration

Beverly Law (UOregon)

Youngryel Ryù (UCB póstdoc) Tara Stiefl (UO student)

Chris Williamson (UO student)

Gretchen Miller (UCB student)

Tom Boden (ORNL)

North American Carbon Program

Peter Thornton (ORNL) Kevin Schaefer (NSIDC)

Daniel Ricciuto (ORNL)

FLUXNET Collaboration

Dennis Baldocchi (UCB)

Rodrigo Vargas (UCB postdoc) Dario Papale (UTuscia)

Markus Reichstein (Máx Plank)

Bob Cook (ORNL)

Dorothea Frank (Max Plank)

National Marine Fisheries Service

Charlotte Ambrose (NOAA)

Pat Stiefer (SEC)

Sonoma County Water Agency

Chris Delanev

The saga continues at: http://bwc.berkeley.edu and http://www.fluxdata.org








Microsoft^{*} Research



































Office of Science